

# Rare Animals of Louisiana




## American Swallow-tailed Kite

*Elanoides forficatus*

**Rarity Rank:** S1S2B/G5

	1	2	3	4	5
State					
Global					
	imperiled		rare		secure



© John Cang www.johncangphoto.com

### Identification:

- A raptor with long, pointed wings and a deeply forked tail
- White head, neck and parts under with black upperparts, wings and tail
- Immatures have white tips on black feathers and fine, dark streaks on the head and breast
- Largest of the North American kites
- Adult length ranges from 51 to 61 cm with a wingspread of 1.3 m

### Habitat:

- Prefers lowland areas particularly in the coastal plain along river systems
- Pines adjacent to swampland particularly attractive to nesters
- Areas with tall trees that can be used for nesting and with open areas to catch prey

### Food habits:

- Insects, small snakes, lizards, frogs, and nestling birds
- Catches and eats most prey while in flight

### Range:


- Breeding Range: northern subspecies from South Carolina to Florida, west to Louisiana; southern subspecies from southeastern Mexico to northern South America
- Winters in South America
- In Louisiana, migrants may be observed statewide

### Reproduction:

- Nests are made of twigs and are up to 50 cm across
- Nests are lined with Spanish Moss and are located near the tops of large trees along rivers or forest openings
- Lays two or three eggs which are shades of white with brown and reddish-brown markings
- Incubation, by male and female, lasts about 28 days, young fledge at five to six weeks

### Threats:

- Vulnerable to harassment by humans and low-flying aircraft while roosting
- Habitat loss caused by a variety of actions including agricultural, urban development, logging, and flood control



Range based on occurrences in Natural Heritage database

# Rare Animals of Louisiana



- Harassment on wintering grounds is a major threat to northern species

## ***Beneficial Management Practices:***

- Limit timber harvest near active nest trees to provide buffer for nesting birds
- Restrict airspace over large roosting areas during breeding season and prevent on ground human disturbance
- Thinning of some timber stands may benefit nesters

## ***LA River Basins:***

Pearl, Pontchartrain, Terrebonne, Atchafalaya, Vermilion-Teche, Mermentau, Calcasieu, Sabine



© Korhnak

Photo by Larry Korhnak

## ***References:***

Meyer, K. D. 1995. Swallow-tailed Kite (*Elanoides forficatus*). In The Birds of North America, No. 138 (A. Poole and F. Gill, eds.). The Academy of Natural Sciences, Philadelphia, and The American Ornithologists' Union, Washington, D.C.

Funding provided by the Louisiana Department of Wildlife and Fisheries and the Barataria-Terrebonne National Estuary Program

For more information, please visit our Web pages at

[www.wlf.louisiana.gov/experience/naturalheritage](http://www.wlf.louisiana.gov/experience/naturalheritage) or 225-765-2811

[www.BTNEP.org](http://www.BTNEP.org) or 1-800-259-0869